

IMPORTANCE OF IRON (Fe) AND COPPER (Cu) FOR CITRUS

Dinesh Kumar¹; Kishore Kumar Das² and Sunil Kumar³

^{1&3} Department of Soil Science and Agricultural Chemistry, ²Department of Horticulture, College of Agriculture, SKRAU, Bikaner -334006

Corresponding Author Email: dkrachiata@gmail.com

Abstract: citrus is a nutrient loving plant and about 15 elements have been known to have important role to play for proper growth and development of citrus. In addition to the major nutrients like N, P, K, Ca, Mg and S, citrus requires micronutrients like Zn, Cu, Mn, Fe, B, Mo etc. Inadequate plant nutrition causes serious disorder in citrus and may eventually lead to decline of the orchard. Iron and copper deficiency leads to serious damage to the citrus production. The importance and role of Iron and copper is described in this manuscript which will be useful to the citrus farming community.

Introduction:

Plant nutrients are classified as macronutrients and micronutrients. Both are essential for plant growth, but macronutrients are needed in large amounts, while micronutrients are needed only in small amounts. The micronutrients are manganese, zinc, iron, copper, boron, molybdenum, chlorine and nickel. When macronutrients are supplied in relatively high proportions compared with micronutrients to stimulate growth of young citrus trees, extreme depletion of micronutrients can develop as a result of marked top growth, and micronutrient deficiencies can appear. Therefore, a balance between macronutrients and micronutrients is needed. Citrus micronutrient deficiencies are most commonly observed in, but not limited to, previously uncultivated soils, shallow soils with high water table, extremely sandy areas and calcareous soils. Since mineral nutrition is a major factor in maximizing yield of high-quality fruit, it is essential to understand the functions of mineral elements, diagnose nutrient deficiencies, and provide needed fertilizers in sufficient amounts at the right time and in the right place. This article describes and discusses iron (Fe) and copper (Cu) deficiencies, functions and recommended practices to alleviate nutritional problems.



Figures 1 and 2 Iron deficiency symptoms.

Iron and Copper Deficiencies

Iron deficiency is common on calcareous soils. A calcareous soil contains a high concentration of calcium carbonate and has a pH of about 8.3. These soils may contain

appreciable amounts of Fe, but it exists in a form that is only slightly available to plants. Iron deficiency can also be attributed to low soil Fe concentration in white sandy areas near lakes or ponds, and in places known locally as “sand-soaked areas.” Iron deficiency can be induced by high phosphorus (P) or accumulation of heavy metals — primarily Cu — in the soil. Iron deficiency is commonly associated with zinc (Zn) and manganese (Mn) deficiencies.

The symptom of Fe deficiency is known as “iron chlorosis” and is called “lime-induced chlorosis” when it occurs on calcareous soils. Deficiency symptoms occur on young leaves, which appear light yellowish to white in color, with the veins greener than the remainder of the leaf (Figure 1). In acute cases, the leaves are reduced in size, fragile, very thin, and can shed early. The trees die back severely on the periphery and especially at the top (Figure 2). Some trees may have a dead top with the lower limbs carrying almost normal foliage. Eventually tree canopy volume decreases and fruit set and yield are reduced. The fruits tend to be small with reduced soluble solids and acidity in the juice. Sometimes only a branch of a tree may be affected or perhaps only a few trees in a grove will be chlorotic. In severe cases, the entire tree is affected. The tree loses some of its leaves, becomes unproductive and dieback results. Iron deficiency is usually associated with high soil alkalinity, but it is also associated with excessive irrigation, prolonged wet soil conditions or poor drainage, and low soil temperature. Some areas displaying Fe chlorosis have been substantially helped or completely cured by careful control of irrigation and drainage. Iron deficiency sometimes occurs where excess salts are present in the soil and in other situations that are not clearly understood.

Copper deficiency is known as “dieback,” “ammoniation” and “exanthema.” These names are derived from the dying back of the twigs, frequent association with heavy application of N (ammonia) and exudates on the surface of the twigs and fruit. Excessive N fertilizer application has been considered for years as a contributing cause of this problem, giving rise to the term “ammoniation.” The cause might be an excessive N/Cu ratio in the plant. This condition is classified as Cu deficiency as opposed to excessive N, and the term “copper deficiency” is preferred in a practical sense because Cu application solves the problem.

The first symptom of Cu deficiency is formation of unusually vigorous, large, dark green foliage with a “bowing up” of the midrib. Twigs are also unusually vigorous, long, soft, angular, frequently “S” shaped, and somewhat drooping. As the deficiency becomes acute and the twigs start to die, some of the weak twigs will bear very small, yellowish-green leaves that drop quickly, leaving the twig defoliated. Fruit symptoms are most pronounced on oranges. Brown-stained areas of hardened gum on the fruit rind may precede the appearance of leaf and twig symptoms. Fruit splitting is common on the trees showing mild symptoms of Cu deficiency, with a part of the splitting beginning at the blossom end in the usual way but some of it occurring across the axis or around the gum-stained areas. In severe cases, dieback of young twigs occurs, and reddish brown droplets of gum cover the twigs. The brown-stained areas on the fruit darken and converge as time progresses and may become almost black by the time the fruit is half grown. Most of this fruit will shed by summer. Insufficient available Cu in the soil is believed to be the primary cause of the

symptoms described. When Cu deficiency is combined with either Zn or Mg deficiency, leaf and twig symptoms may not be noticed due to reduction in growth, but the typical fruit symptoms will still show. Therefore, fruit symptoms are the most reliable indicator in routine observation.

Copper deficiency is more of a problem on newly planted flat woods soils compared with deep sandy soils of the Central Florida ridge. Copper deficiency can be a controlling factor in fruit production, and acute Cu deficiency may take trees entirely out of production.

Functions of Iron

- * Catalyzes the production of chlorophyll
- * Involved in some respiratory and photosynthetic enzyme systems
- * Involved in the reduction of nitrates and sulfates

Functions of Copper

- * Part of several enzyme systems
- * Has a role in photosynthesis and chlorophyll formation
- * May have an important function in root metabolism
- * Regulates several biochemical processes within the plant
- * Important in the utilization of proteins in growth processes
- * Involved in oxidation-reduction reactions

Table1. Guidelines for interpretation of orange tree leaf Fe and Cu (mg/kg or ppm) analysis based on 4- to 6-month-old spring flush leaves from non-fruiting twigs collected in July–September**

Element	Deficient	Low	Optimum	High	Excess
Fe	<35	35 – 59	60 – 120	121 – 200	>200
Cu	<3	3 – 4	5 – 16	17 – 20	>20

**Milligrams/kilograms or parts per million

Fertilizer Sources and Formulations For Soil Application

Iron

- *Iron oxy-sulfate
- *Iron EDTA and HEDTA Iron DTPA
- *Iron EDDHA Iron humate

Copper

- *Copper sulfate

Recommendations

Iron is not easy to remedy iron chlorosis of citrus trees on susceptible rootstocks (Swingle citrumelo, trifoliolate orange) planted on calcareous soils. Iron fertilizer formulations are available that can correct chlorosis, but the required application rate and frequency make the treatment expensive. Inorganic sources of Fe such as ferrous sulfate (FeSO₄) or ferric sulfate [Fe₂(SO₄)₃] are not effective. Iron chlorosis should be addressed through soil application of Fe chelates. Chelates are superior Fe sources for plants because they supply sufficient Fe at much lower rates compared with inorganic Fe sources. The most popular synthetic organically chelated forms of Fe include Fe-EDTA, Fe-HEDTA, Fe-DTPA, and Fe-EDDHA. The effectiveness of these fertilizers varies greatly depending on soil pH.

Organically-complexed Fe exists in byproducts such as wastewater residuals

(biosolids) or certain drinking water treatment residuals (Fe-humates). Biosolids are potentially useful because they contain high concentrations of Fe in a complex form that does not readily precipitate. Research with Fe-humates applied to citrus trees showed that moderate Fe deficiency could be corrected for relatively low cost. Foliar application of FeSO₄ or Fe chelates has not proven satisfactory on citrus trees because of poor translocation within the leaf. The use of foliar sprays also increases the possibility of fruit and/or leaf burn. For these reasons, foliar application of Fe is not recommended to correct Fe chlorosis of citrus. Citrus rootstocks vary widely in their ability to overcome low Fe stress. Swingle citrumelo and trifoliolate orange are among the most sensitive rootstocks to Fe chlorosis. The easiest way to avoid alkaline-induced Fe chlorosis in citrus trees that are planted on calcareous soils is to use tolerant rootstocks.

Copper Foliar sprays or soil applications of Cu fertilizer can prevent or cure Cu deficiency. Spraying a solution containing 2 to 3 pounds per acre of elemental Cu applied during flowering commonly results in near-immediate recovery followed by a normal fruit set. Foliar sprays are useful emergency treatments when symptoms of Cu deficiency are first observed.

Some central ridge and east coast grove soils contain 300 to 400 pounds of Cu/acre in the top 6 inches of soil. A moderate routine disease control spray program contributes an additional 3 to 4 pounds Cu/acre/year, so no soil application of Cu fertilizer is needed in this situation. Soil pH below 5.5 can solubilize enough soil Cu to be toxic to citrus trees. As little as 1 ppm Cu in the soil solution can kill roots. Maintaining soil pH close to neutral is recommended to reduce the potential for Cu toxicity. Some rootstocks (e.g. Swingle citrumelo) that perform poorly in high pH soils are also sensitive to high soil Cu. When replanting on old grove sites with low soil pH, the potential for toxicity is high. While the emphasis with old grove soils is on managing high soil Cu, most previously non-cultivated Florida sandy soils are deficient in Cu. If a grove is developed on a virgin sandy soil, Cu should be included in the fertilizer blend at 2.5 percent of the nitrogen rate.

References

- Reuther, walter; smith, paul f (1993)** effects of high copper content of sandy soil on growth of citrus seedlings *Soil Science*, **75:3**
- F. Behlau, J. Belasque Jr.J.H., Graham, R.P. Leite Jr. (2010)** Effect of frequency of copper applications on control of citrus canker and the yield of young bearing sweet orange trees. *Crop Protection* **29:3**
- Longo, A.Benintende (2002)** Mineral nutrition of Citrus trees. Effect of copper deficiency single or associated on free amino acid content of leaves [Citrus aurantium L.] P.G. (Catania Univ. (Italy). Dipartimento di Scienze Agronomiche, Agrochimiche e delle Produzionin
- Abdolhossein Aboutalebi (2013)** Effects of nitrogen and iron on Sweet lime (Citrus limmetta) fruit quantity and quality in calcareous soils. *Journal of Novel Applied Sciences* **2: 8**
- SE El-Kassas (2010)** Effect of iron nutrition on the growth, yield, fruit quality, and leaf composition of seeded Balady lime tress grown on sandy calcareous soils. *Journal of plant Nutrition* vol **7: 1-5**