



ROLE OF REMOTE SENSING IN PRECISION AGRICULTURE

*Basavaraj Patil and Chetan, H. T.

University of Agricultural Sciences, GKVK, Bengaluru-65

Email of corresponding author - bspatil4504@gmail.com

Introduction: Practices in developed countries have tended to support greater energy inputs using large machineries and increased applications of chemicals and fertilizers. While these practices have negative societal and environmental implications such as soil erosion and salinization, soil fertility, compaction of subsoils and soil/water pollution, they have generally supported the food and fiber needs of a rapidly growing human population. A paradigm shift toward a new production method that ensures safe and sustainable agriculture is needed. Across the globe, Precision Agriculture (PA) is changing the way people farm as it offers a myriad of potential benefits in profitability, productivity, sustainability, crop quality, environmental protection, on-farm quality of life, food safety and rural economic development. PA is an innovative, integrated and internationally standardized approach aiming to increase the efficiency of resource use and to reduce the uncertainty of decisions required to manage variability on farms. PA has been hailed as one of the most scientific and modern approaches to production agriculture in the 21st century, as it epitomizes a better balance between reliance on traditional knowledge and information and management-intensive technologies. At present, there is an increasing commitment to reduce reliance on excessive chemical inputs in agriculture. Numerous technologies have been applied to make agricultural products safer and to lower their adverse impacts on the environment, a goal that is consistent with sustainable agriculture. PA has emerged as a valuable component of the framework to achieve this goal (Zhang *et al.*, 2002).

Precision Agriculture (PA)

PA is an integrated, information- and production based farming system that is designed to increase long term, site-specific and whole farm production efficiency, productivity and profitability while avoiding the undesirable effects of excess chemical loading to the environment or productivity loss due to insufficient input application. The inference is that better decision making will provide a wide range of benefits in economic, environmental and social aspects that may or may not be known or measurable at present (Auernhammer, 2001). Worldwide, investments in research and technology development on PA have considerably increased during the past decade (Schellberg *et al.*, 2008).

Importance of precision Agriculture

The importance of PA has been widely recognized as a key contributor in crop production technology around the globe, but so far, this technology is only becoming practicable on large farms. PA is based on innovative systems approach and these new systems approach depends on a combination of fundamental technologies such as Geographic Information System (GIS), Global Positioning System (GPS), computer modeling, ground based/airborne/satellite remote sensing, variable rate technology and advanced information processing for timely in-season and between season crop management. Protocols for PA implementation can be encapsulated in three general steps: (1) Gathering information about variability, (2) Processing and analyzing

information to assess the significance of variability and (3) Implementing change in the management of inputs. PA subscribes to a cyclic process that is typified by a system that gets smarter every year a farm operator uses it. Taken together, this implies that the three general steps may each require a timeframe ranging from months to years so as to collectively generate a stable and workable structure, but yet be subjected to further amendments and refinements. The importance of PA evolved during the era of agricultural mechanization in the 20th century where there was strong economic pressure to treat large fields with uniform agronomic practices. PA provides a means to automate Site-Specific Management (SSM) using information technology, thereby making SSM practical in commercial agriculture (Auernhammer, 2001).


The PA database generally includes (Venkataratnam, 2001):

Crop information such as growth stage, health, nutrient requirement

- Soil physical and chemical properties, depth, texture, nutrient status, salinity and toxicity, soil temperature, productivity potential
- Microclimatic data (seasonal and daily) such as canopy temperature, wind direction and speed, humidity
- Surface and sub surface drainage conditions
- Irrigation facilities, water availability and planning of other inputs

A PA system that harnesses recent advances in sensor technology can play a crucial role toward an intelligent crop production system. Specifically, remote sensing technology that allows non-destructive acquisition of information about the Earth's surface can facilitate the implementation of PA. For example current crop status (including maturity period) and crop stresses such as nutrient and water stress, disease, pest and weed infestations can be discerned by means of remote sensing instruments such as cameras, laser scanners, linear arrays and area arrays, without actually being in contact with them. Information gathered via different sensors and referenced using a GPS can be integrated to create field management strategies for chemical application, cultivation and harvest. This review highlights on remote sensing technology and describes how it can be used as an effective tool in Precision Agriculture.

Fig. 1: The visible region of the spectrum ranges from about 0.4-0.7 μm



Remote sensing technology: Remote Sensing (RS) is the science of obtaining and interpreting information from a distance, using sensors that are not in physical contact with the object being observed (Jensen, 1996). The science of remote sensing

includes aerial, satellite and spacecraft observations of the surfaces and atmospheres of the planets in our solar system, while the Earth is noticeably the most frequent target of study. RS is

usually restricted to methods that detect and measure electromagnetic energy including visible and non-visible radiation that interact with surface materials and the atmosphere (Fig. 1). RS and GIS technologies have been of great use to planners in planning for efficient use of natural resources at national, state and district levels. Application of these technologies in the management of natural resources are increasing rapidly due to great strides made in space-borne RS satellites in terms of spatial, temporal, spectral and radiometric resolutions (Venkataratnam, 2001).

Remote sensing has several unique advantages (Jensen, 1996):

- RS technology is well-known as a non destructive method to collect information about earth features
- RS data may be obtained systematically over very large geographical areas rather than just single point observations
- RS data can reveal information about places that are inaccessible to human exploration
- The systematic (raster) data collection in RS can remove sampling bias
- RS can provide fundamental biophysical information that can be used in other sciences
- RS is independent from the data produced elsewhere, in comparison with the other mapping sciences such as cartography or GIS

The history of modern RS began when black and white photographs of the landscape were first taken from the air. The first utilization of aerial photographs appeared for military purposes. Eventually, the Agriculture Stabilization and Conservation Service adopted the technology and began collecting indexed photographs of the landscape for agricultural purposes. RS, today, incorporates new technologies that provide increasingly efficient, complete, accurate and timely information.

How can remotely-sensed data be used in agriculture?

Today, RS is potentially a practical management tool for site-specific crop management. Currently, there is a wide range of satellite data that varies in (i) technique (active/ passive, radiometer/ scatterometer), (ii) spatial resolution from submeter to kilometers (iii) spectral range, and (iv) viewing geometry. The full commercial availability of very high resolution satellite data has opened up a number of new opportunities for the use of Earth Observation (EO) data. Today, we can perform many applications with EO data that in only the recent past were exclusive to manpower investigation and *in situ* surveys, which was time consuming and hard-sledding, despite the geographic limitations of such data and techniques. Satellite imagery can be acquired over any area globally, in a time frame and at a given price. At present, higher resolution satellite imagery overcomes previous constraints and permits the use of such data as a quick and easy tool for territorial management, including agricultural analysis, statistics and subsidy control. QuickBird is currently the satellite with the highest resolution which is available for agricultural and civilian uses.

New RS multispectral and hyperspectral sensors are swiftly generating vast amounts of data in a cost-effective manner and at higher spatial and spectral resolutions. Hyperspectral and multispectral images, consisting of reflectance from the visible, near infrared and mid-infrared regions of the electromagnetic spectrum, can be interpreted in terms of physical parameters (such as crop cover, crop health and soil moisture) and are useful for operations such as stress mapping, fertilization and pesticide application and irrigation management (Singh *et al.*, 2007). Nutrient

contents of different crops such as wheat, paddy, sorghum, corn, broccoli, citrus, grape, and apple have also been assessed using hyperspectral and multispectral RS data. Interpretation of RS data is often aided by specialized techniques such as geostatistics, image analysis and classification, and artificial intelligence.

RS techniques also play an important role in assessing crop condition and yield forecasting, acreage estimates of specific crops, detection of crop pests and diseases, disaster location and mapping, wild life management, water supply information and management, weather forecasting, rangeland management, and livestock surveys.

Some disease and insect pests of crops may be monitored by remote sensing. Riedell *et al.* (2004) introduced remote sensing technology as an effective and inexpensive method to identify pest-infested and diseased plants. They used remote sensing techniques to detect specific insect pests and to distinguish between insect and disease damage on oat. Results suggested that canopy characteristics and spectral reflectance differences between insect infestation damage and disease infection damage can be measured in oat crop canopies by remote sensing but that these differences may not be consistent from one growing season to the next.

RS applications in agriculture have progressed to a stage where information from RS imagery is being used for a number of policy level decisions related to food security, poverty alleviation and sustainable development. Decision on buffer stock of food grains could be based on pre-harvest crop acreage and production estimates while the ground water potential maps serve as a major source of information in ensuring drinking water and other needs in rain-fed and less favorable areas. Nationwide land use, land cover, soil and wasteland mapping have helped in expansion and intensification of agricultural activities and also in identification of land capability classes and crop suitability indices (Venkataratnam, 2001).

References:

- Auernhammer, H., (2001).** Precision farming-the environmental challenge. *Comput. Elect. Agric.*, **30**: 41-33.
- Jensen, J. R., (1996).** Remote sensing of the environment: *An Earth Resource Perspective*. 3th Edn., Prentice Hall, USA, pp: 1-28.
- Riedell, W.E., S.L. Osborne and L.S. Hesler, (2004).** Insect pest and disease detection using remote sensing techniques. Proceedings of 7th International Conference on Precision Agriculture. Minneapolis, MN USA.
- Schellberg, J., M.J. Hill, R. Gerhards, M. Rothmund and M. Braun, (2008).** Precision agriculture on grassland: Applications, perspectives and constraints. *Eur. J. Agron.*, **29**: 59-71.
- Singh, D., Sao, R. and Singh, K.P. (2007).** A remote sensing assessment of pest infestation on sorghum. *Adv. Space Res.*, **39**: 155-163.
- Venkataratnam, L., (2001).** Remote sensing and GIS in agricultural resources management. Proceedings of the 1st National Conference on Agro-Informatics, June 3-4, Dharwad, India, pp: 20-29.
- Zhang, N., M. Wang and N. Wang, (2002).** Precision agriculture: A worldwide overview. *Comput. Elect. Agric.*, **36**: 113-132.